

271570

STIC-ILL

From: Zeman, Mary  
Sent: Thursday, November 18, 1999 3:00 PM  
To: STIC-ILL  
Subject: ILL Reference Request 09/095,385

Scientific and Technical  
Information Center

NOV 23 RECD

PAT. & T.M. OFFICE

Please send a copy of the following references.

Thank you  
Mary Zeman  
305 7133  
Examiner, 1643  
8D17

Current Topics in Microbiology and Immunology Dec 1999 Vol 236 pages 93-111  
ISSN: 0070-217X

Genetic Engineering and Biotechnologist 1995 Vol 15 no 4 pages 249-259  
ISSN: 0959-020X

Infection and Immunity 1993 (dec) vol 61 no 12 pages 5279-5285  
ISSN: 0019-9567

Advances in Experimental Medicine and Biology 1991 vol 310 pages 183-185  
ISSN: 0065-2598

Journal of Cell Biology May 1989 108 5 1673-1685  
ISSN: 0021-9525

NOS

99109235

991122B171

reh  
11/23  
Vao  
NOS  
in use

# Antibody-Mediated Protection of Mucosal Surfaces

B. CORTHEY<sup>1</sup> and J.-P. KRAEHNBUHL<sup>2</sup>

|       |  |     |
|-------|--|-----|
| 1     | Introduction   | 93  |
| 2     | Humoral Immune Responses in Mucosal Tissues                  | 94  |
| 3     | Maturation of B Cells within Plasma Cells in Mucosal Tissues | 96  |
| 4     | Trans epithelial Transport of Mucosal Immunoglobulins        | 97  |
| 5     | Antibodies in Mucosal and Glandular Secretions               | 99  |
| 5.1   | Molecular Structure of Immunoglobulin A (IgA)                | 99  |
| 5.1.1 | Polypeptide Composition of Secretory (s) IgA                 | 100 |
| 5.1.2 | Glycosylation of sIgA  | 101 |
| 5.1.3 | Regulation of SC and $\alpha$ Chain Gene Expression          | 102 |
| 5.2   | Functions of sIgA Antibodies                                 | 103 |
| 5.2.1 | Immune Exclusion by Antigen Cross-Linking                    | 103 |
| 5.2.2 | Interaction with Mucus                                       | 104 |
| 5.2.3 | Neutralization and Opsonization                              | 104 |
| 5.2.4 | Intracellular Neutralization and Antigen Clearance           | 105 |
|       | References   | 106 |

## 1 Introduction

Mucosal surfaces of the oral cavity, the digestive and urogenital tracts, and the airways are protected against environmental pathogens by innate and adaptive immune defense mechanisms. Innate defense involves physical, chemical and cellular factors. Entrapment of pathogens in mucus facilitates their clearance by peristalsis in the gut and ciliary movement in the airways. The longitudinal flow of fluids across the epithelial layer mediated by chloride channels helps to flush away microorganisms and prevent their attachment to the epithelial cell surface. The apical cell surface-associated glycocalyx, which consists of a dense network of

<sup>1</sup>Division of Immunology and Allergology, Centre Hospitalier Universitaire Vaudois, 1011 Lausanne, Switzerland

<sup>2</sup>Swiss Institute for Cancer Research and Institute of Biochemistry, University of Lausanne, 1066 Epalinges, Switzerland

glycoproteins, also limits the access of pathogens to the epithelial surface. Chemical factors (lysozyme, lactoferrin, peroxidase and defensins) secreted by specialized epithelial cells, such as the Paneth cells in the crypts of the small intestine, gastric acid and intestinal hydrolases (proteases, lipases, nucleases) constitute an efficient defense mechanism.

Host leukocytes present or recruited into mucosal tissues following microbial-epithelial cell interactions actively participate in innate defense. The recruited cells include monocytes, macrophages, neutrophils and eosinophils. These cells preferentially express Fc $\alpha$  receptors for immunoglobulin A (IgA) antibodies in both mice (HAYAMI et al. 1997) and humans (MONTEIRO et al. 1990) in mucosal tissues. These Fc $\alpha$  receptor-bearing cells may help to clear microorganisms, following opsonization by secretory IgA (sIgA) antibodies, facilitate uptake of antigen-antibody complexes by professional antigen-presenting cells and modulate immune functions by controlling the release of specific cytokines. Therefore innate immunity may not only provide rapid anti-microbial defense, but may also determine how pathogens activate the adaptive mucosal immune system and control the nature of the immune response (FEARON and LOCKSLEY 1996; BENDELAC and FEARON 1997). Adaptive immunity requires that the antigens cross the epithelium to reach the underlying organized lymphoid tissues where antigen presentation and priming of T and B lymphocytes can occur.

## 2 Humoral Immune Responses in Mucosal Tissues

sIgA antibodies play a major role in the protection of mucosal surfaces which is reflected by the very large quantity of antibodies produced and transported into secretions each day (~10g). Following stimulation by antigens, naive B cells in organized mucosal lymphoid tissues (MALT) of the gut, the airways and the oropharyngeal cavity move to the germinal center where they proliferate clonally. During clonal expansion, B cells undergo affinity maturation, first, by somatic hypermutation which generates variability in B-cell receptors and, second, by selection of those with highest affinity for the antigen. Selection of cells bearing these mutated receptors by the antigen occurs on the surface of the follicular dendritic cells a process which rescues cells expressing high affinity Ig receptors from apoptosis (for review, see LIU et al. 1992).

In MALT germinal centers, B lymphocytes undergo isotype switch and differentiate further into B cells that express IgA receptors. (CEBRA et al. 1991). MALT CD4<sup>+</sup> T cells have been shown to promote IgA isotype switch of IgM-bearing B cells (KAWANISHI et al. 1983). Cytokines produced by activated Th2 helper CD4<sup>+</sup> T cells, including IL-5, IL-10 and transforming growth factor beta (TGF- $\beta$ ) play a major role in triggering switch, but the precise molecular mechanism that mediates the recombination event has not yet been fully elucidated (STROBER and EHRHARDT 1994). Ligation of co-stimulatory molecules, such as

CD40, provides an important signal for switch induction (DEFRANCE et al. 1992). Bacterial lipopolysaccharide (LPS) has been shown to stimulate expression of the recombination machinery in pre-B lymphocytes (LI et al. 1996) and mucosal adjuvants, including cholera toxin and *E. coli* heat-labile toxin, are known to facilitate switch (LYCKE and STROBER 1989).

Subsequently, B lymphocytes differentiate into effector or memory cells, following contact with T helper lymphocytes and CD40-CD40 ligand interactions (LIU et al. 1991). In MALT, stimulated B and T cells acquire a mucosal homing program. The effector and memory lymphocytes lose their adhesion to stromal cells, leave organized MALT structures and enter the blood stream via the lymph. Depending on the mucosal sites at which priming takes place, different homing receptors will be expressed by B lymphocytes. Virtually all IgA- and even IgG-antibody-secreting cells detected after peroral and rectal immunization expressed  $\alpha 4\beta 7$  integrin receptors, with only a minor fraction of these cells expressed the peripheral L-selectin receptor. In contrast, circulating B cells, induced by intranasal immunization, co-express L-selectin and  $\alpha 4\beta 7$  receptors. This may explain the compartmentalization of mucosal immune responses initiated in the upper vs the lower aerodigestive tract (QUIDDING-JARBRINK et al. 1997).

Effector and memory B cells are able to home to distant mucosal tissues or return to MALT structures (ROTT et al. 1997 and, for review, BUTCHER and PICKER 1996) (Fig. 1). The lymphocytes expressing mucosal  $\alpha 4\beta 7$  homing receptors interact with flat post-capillary venule endothelial cells bearing mucosal addressins on their luminal surface (for review, see BUTCHER and PICKER 1996). Antigen receptors (surface Igs) do not participate in the selectivity of lymphocyte binding to the vascular bed. It has been suggested that antigen-specific plasmablasts become locally enriched in mucosal sites via retention at sites of antigen deposition (BUTCHER and PICKER 1996).

The mucosal addressin MadCam-1 (mucosal addressin cell adhesion molecule) is preferentially expressed in human and mouse intestinal flat post-capillary venules of the lamina propria and high endothelial venules of organized MALT, but not in other mucosal tissues (BRISKIN et al. 1997). MadCam-1, in mice, is also found associated with the endothelial cells of mammary-gland post-capillary venules. The vascular addressins mediating selective binding of lymphocytes in the airways and the genital tract have yet not been identified. Extravasation of lymphocytes requires the action of chemokines (for review, see BUTCHER and PICKER 1996). B lymphocytes express a specific chemokine receptor, which when knocked out prevents lymphocytes from reaching mucosal tissues (FÖRSTER et al. 1996). The chemokine that attracts B cells has recently been identified (GUNN et al. 1998; LEGLER et al. 1998), but it is not known whether B cell-specific chemokines are involved in the recruitment of B cells in mucosal tissues.




Fig. 1. In stratified epithelia (tonsil, vagina), antigen sampling is accomplished by migratory dendritic cells that transport antigens to distant lymph nodes or to local organized mucosa-associated lymphoid tissues (O-MALT). In simple epithelia (intestine, bronchi), membranous (M) cells selectively take up the antigens and deliver them to underlying local O-MALT. Following antigen stimulation in lymphoid tissues, effector lymphocytes enter the submucosal lymphatics enter the blood circulation and establish themselves in distant MALT and glandular sites. Migration to effector sites depends on the specific interaction between homing receptor on the lymphocytes and addressin on the vascular bed bathing the epithelia. B-cell differentiation into antibody-producing cells leads to the local production of immunoglobulin A (IgA) which, upon association with the polymeric immunoglobulin receptor (pIgR), are transported across the epithelial layer and serve to neutralize pathogenic agents

### 3 Maturation of B Cells within Plasma Cells in Mucosal Tissues

Locally, in the lamina propria, effector B lymphocytes differentiate into antibody-secreting plasma cells, this process being regulated by T lymphocyte-, and epithelial, cell-derived cytokines (Fig. 2). In the intestinal mucosa, the number of plasma cells producing IgA exceeds those producing all other Ig isotypes. Maturation of IgA-bearing B cells into plasma cells is triggered by T cell-derived IL-5 (MATSUMOTO et al. 1989) and epithelial IL-6 (MCGHEE et al. 1991). In IL-6-deficient mice, a reduced number of IgA-producing plasma cells has been observed in the respiratory tract, and targeting IL-6 DNA into bronchial epithelial cells restored maturation of IgA B cells into plasma cells (RAMSAY et al. 1994). This result, however, has not been confirmed within the digestive tract, suggesting that IL-6 is probably not the only cytokine involved in B cell maturation (BROMANDER et al. 1996). In the mucosal environment, all plasma cells, irrespective of their Ig isotype, express J chain, the small polypeptide required for IgA polymerization (BJERKE and BRANDTZAEG 1990).




Fig. 2. Role of cytokines in the immunoglobulin A (IgA) response and biosynthesis. Mature (antigen responsive) B lymphocytes expressing membrane IgM switches to surface IgA production, under the control of Th2 cytokines, including transforming growth factor beta (TGF- $\beta$ ), interleukin (IL)-4, IL-5 and IL-10. This happens most likely in the germinal centers of organized mucosa-associated lymphoid tissues (O-MALT). IgA<sup>+</sup> cells eventually leave the O-MALT and migrate to distant mucosal and glandular sites to undergo final differentiation. Enterocyte-derived IL-6 and Th2-released IL-5 and TGF- $\beta$  induce secretion of soluble IgA oligomers through concomitant induction of J chain expression. Following binding to polymeric immunoglobulin receptor (pIgR), J chain-containing IgA polymers are selectively transported across the epithelium by transcytotic vesicles which, following fusion with the apical membrane, release sIgA complexes comprising IgA and the clipped form of pIgR called secretory component (SC). The possible sites of action of IgA/sIgA are listed on the right and discussed case by case within the text.

#### 4 Transepithelial Transport of Mucosal Immunoglobulins

The prevention of microbial infection requires that the humoral immune effectors gain access to the luminal compartment to block adhesion and invasion of environmental pathogens. Simple epithelia are sealed at their apex by tight junctions that prevent the lateral diffusion of antibodies between cells. Thus, translocation of antibodies from their site of synthesis in the lamina propria to the luminal compartments necessitates an efficient transepithelial transport machinery (Fig. 2). The pathway followed by polymeric Igs (IgA and IgM) across epithelial cells has been extensively studied (SOLARI and KRAEHNBUHL 1984; MOSTOV and DEITCHER 1986, BARROSO and SZTUL 1994; NATVIG et al. 1997; WEIMBS et al. 1997), and the receptor that mediates transcytosis has been well characterized (EIFFERT et al. 1984; KRAJCI et al. 1989; SCHAEFER et al. 1991; COYNE et al. 1994).

The polymeric immunoglobulin receptor (pIgR) is expressed at the basolateral surface of a variety of epithelial cells, including small and large intestinal, nasal, tracheal and bronchial, and cervical and uterine epithelial cells of all mammals studied so far (BRANDTZAEG 1994). The receptor comprises five extracellular domains of 110–120 residues which share homology with V $\kappa$  and V $\text{H}$  Ig domains, a 23-amino acid long membrane-spanning segment, and a cytoplasmic tail of about 100 amino acids (MOSTOV et al. 1984). Human, rat, mouse, rabbit and bovine receptors show extensive homology, particularly in the first domain and in the cytoplasmic tail. Following polymeric IgA binding and internalization into the recycling basolateral endosomal compartment, the complex is sorted into transcytotic vesicles that reach the apical endosomal compartment (SONG et al. 1994), where the ligand–receptor complex recycles transiently (APODACA et al. 1994).

The presence of the J chain in polymeric IgA accounts for the selective discrimination of polymeric vs monomeric IgA (BRANDTZAEG and PRYDZ 1984). At the apical surface, the poly-Ig receptor is proteolytically cleaved, and the extracellular fragment known as secretory component (SC) is released, whether bound or not to its polymeric IgA ligand. Rabbits express a spliced version of the poly-Ig receptor lacking domains 2 and 3 (DEITCHER and MOSTOV 1986) which is fully competent for transport (SOLARI and KRAEHNBUHL 1987) and generates SC non-covalently bound to IgA (FRUTIGER et al. 1987). The information necessary for the complex trafficking of the pIgR in epithelial cells resides within its cytoplasmic tail. The sequences that control basolateral sorting and endocytosis (MOSTOV et al. 1992), which are highly conserved among species (BANTING et al. 1989), are flanked by serine residues that undergo phosphorylation once the receptor is terminally glycosylated in the Golgi apparatus (HIRT et al. 1993). Transcytosis of the poly-Ig receptor is stimulated by binding of polymeric IgA (SONG et al. 1994). Endocytosis and transcytosis are regulated by the trimeric G proteins, protein kinase C and calmodulin (CHAPIN et al. 1996). In rat liver, the asialoglycoprotein receptor (ASGP-R) has also been shown to participate in IgA endocytosis (SCHIFF et al. 1986). In humans, transport of IgA from blood to bile is minimal, due to the lack of pIgRs in hepatocytes.

IgG antibodies are also found in secretions and it has been proposed that they cross the epithelial barrier by transudation. Recently, however, it has been shown that small-intestinal enterocytes express the neonatal Fc receptor (ISRAEL et al. 1997). In newborn rodents, the receptor mediates transport of maternal milk IgG antibodies from the lumen to the lamina propria. Binding of IgG antibodies in the luminal compartment is facilitated by low pH and release in the interstitium by neutral pH (RODEWALD and KRAEHNBUHL 1984). Transport in the opposite direction requires that IgG antibodies are able to reach an acidic compartment in order to bind to Fc receptors and escape to the lysosomal pathway. In the yolk sac, IgGs are taken up by fluid-phase endocytosis and accumulate in an acidic endosomal compartment where binding to neonatal Fc receptors takes place (ROBERTS et al. 1990). Whether the neonatal Fc-receptor mediates IgG transudation in other mucosal epithelia remains to be established.

## 5 Antibodies in Mucosal and Glandular Secretions

In human, roughly 60% of all Igs produced are IgAs, of which about half are selectively transported into external secretions (MESTECKY and MCGHEE 1987; KERR 1990). The majority of sIgA in humans is derived from local synthesis, and not from the circulation. Humans have two subclasses of monomeric and polymeric IgA: IgA1 and IgA2 (KAWAMURA et al. 1992), which differ in their hinge region and in their carbohydrate composition (WOLD et al. 1995). IgA1, but not IgA2, can be cleaved by IgA-specific proteases produced by a number of bacteria (PLAUT 1988). The protease recognizes a distinct sequence in the hinge region of IgA1 antibodies and renders the antibodies less efficient in cross-linking microorganisms. SC does not protect against cleavage (PLAUT et al. 1985). The proportion of IgA1 and IgA2 varies in individual secretions. IgA1-producing cells predominate in most mucosal tissues and glands, including the tonsils, the stomach, the duodenum, the mammary gland and the respiratory tract, while IgA2-secreting cells populate, preferentially, the large intestine and the female genital tract. Antibodies specific for protein antigens are found predominantly in the IgA1 subclass, whereas bacterial LPS and carbohydrate antigens lead preferentially to IgA2 antibody production. In IgA-deficient individuals and animals, a compensatory increase of sIgM in secretions has been reported (PLEBANI et al. 1983), and experimental IgA-deficient mice challenged with *H. felis* and cholera toxin have much higher titers of IgM antibodies than wild-type animals (NEDRUD et al. 1996).

IgE are also locally produced by plasma cells. They play a major role in the lamina propria where they can interact and activate local mast cells. IgE can mediate protection against certain parasites, as demonstrated in response to the feeding of larval ticks (MATSUDA et al. 1990). Recruitment of leukocytes at sites of IgE-dependent immune response by tumor necrosis factor alpha (TNF- $\alpha$ ) released by mast cells could be a critical component of natural immunity to infection of bacteria and other pathogens. It has been reported that not only Th2 cells, but also mast cells and the B cells that mediate IgE responses express the same chemokine receptor, the eotaxin receptor CCR3 (SALLUSTO et al. 1997), which facilitates their recruitment at mucosal sites. Due to their strategic anatomical distribution near surfaces exposed to the external environment and their ability to release mediators of inflammation, mast cells are well suited to function as sentinels of mucosal immunity.

### 5.1 Molecular Structure of Immunoglobulin A (IgA)

Secretory sIgA has unique features that make this antibody particularly well suited to protect mucosal surfaces. Its polymeric structure enhances its avidity for antigens and pathogens, its distinct carbohydrate composition contributes to its



binding to mucus components and also to its resistance to proteolysis (KERR 1990; MESTECKY et al. 1991). Secretory IgA consists of at least two monomeric IgA units and two additional polypeptide chains, a J chain and SC (Fig. 3). These four polypeptides are produced by two distinct cell types. The Ig heavy and the light chains and the J chain are synthesized and assembled by plasma cells, while SC, which corresponds to the five extracellular domains of the poly-Ig receptor, is contributed to by epithelial cells of mucus membranes and exocrine glands.

### 5.1.1 Polypeptide Composition of Secretory (s) IgA

Mammalian Ig  $\alpha$  heavy chain consists of three constant-region domains (C $\alpha$ 1–C $\alpha$ 3), composed of seven  $\beta$  strands in a four–three configuration with intervening loops to create a  $\beta$ -barrel conformation (HUNKAPILLER and HOOD 1989). In humans, non-human primates and rabbits, multiple  $\alpha$ -chain isotypes have been described (SPIEKER-POLET et al. 1993), in contrast to all other species, where only one IgA isotype is found. In human, two subclasses of IgA have been described: IgA1 and IgA2. The major difference between these two subclasses lies in the hinge region, lacking a 13-amino acid stretch in the IgA2 molecule. In addition, the IgA2




Fig. 3. Schematic representation of a dimeric secretory immunoglobulin A (sIgA). Two IgA monomers are depicted in a tail-to-tail arrangement, with J chain covalently linked to one monomer through two disulfide bridges. Secretory component (SC) is made of five Ig-like domains and corresponds to the extracellular portion of polymeric immunoglobulin receptor (pIgR). Domain I contains the information for initial anchoring to oligomeric IgA, while two cysteine residues situated in the IgA  $\alpha$  chain and the domain 5 of SC form a covalent bond in several species. Sugar moieties most likely contributing to the stability and solubility of sIgA are also drawn, yet not on scale

subclass comprises two allotypes: IgA2 m(1) and IgA2 m(2), the major structural differences of which lie in the arrangement of the  $\alpha$  and light chain interchain disulfide bridges. The C $\alpha$ 3 domain displays, at its C terminus, an 18-amino acid tail that, during biosynthesis of IgA, associates with J chain.

J chain is a 15-kDa glycoprotein, covalently linked to polymeric IgA or IgM before secretion by plasma cells. The role of J chain in the polymerization process remains controversial (CATTANEO and NEUBERGER 1987; DAVIS et al. 1989). IgA dimer formation occurs in J chain-deficient mice, but the IgA dimer/monomer ratio in serum is perturbed (HENDRICKSON et al. 1995); hepatic IgA transport is impaired, whereas intestinal, breast milk and nasal IgA levels compare with wild-type animals (HENDRICKSON et al. 1996). J chain-deficient IgA found in secretions, however, is not associated with SC. These data confirm the essential role of J chain in interaction with the poly-Ig receptor (BRANDTZAEG and PRYDZ 1984).

SC represents the extracellular portion of the poly-Ig receptor. The three-dimensional structure of SC has not been elucidated, but molecular modeling of rabbit domains 1 and 2 indicates that both domains consist of the typical nine Ig $\beta$  strands (A, B, C, C', C, D, E, F, G) connected by loops of variable size (COYNE et al. 1994; CORTHESEY et al. 1996). Each domain contains an internal disulfide bond, characteristic of Ig homology units, which apparently links strand B with strand F. In human sIgA, amino acids 14–38 in the N-terminus domain 1 of SC constitute the IgA binding epitope (BAKOS et al. 1991). Cysteine 467 in the distal fifth domain participates in disulfide bridge formation with cysteine 311 of one C $\alpha$ 2 domain of dimeric IgA. It is not known in which cell organelle covalent disulfide linkages are established during transcytosis, but the process is rapid and spontaneous, as reflected in *in vitro* reconstitution experiments (RINDISBACHER et al. 1995). SC delays IgA degradation by intestinal proteases, but does not alter the affinity for the antigen (LÜLLAU et al. 1996).

### 5.1.2 Glycosylation of sIgA

The glycosylation pattern of sIgA is complex and highly heterogeneous (ENDO et al. 1994; WOLD et al. 1994). The 13-amino acid stretch in the IgA1 hinge region contains five O-linked glycans, recognized by the lectin Jacalin (ROQUE-BARREIRA and CAMPOS-NETO 1985), which presumably enhances the flexibility between Fc and F(ab')<sub>2</sub> fragments. The presence of this extra sequence in IgA1 molecules is responsible for the sensitivity of the antibody to bacterial specific proteases as discussed above. Glycosylation of the  $\alpha$  chain is essential for intracellular stability and normal secretion of IgA (TAYLOR and WALL 1988). In contrast, absence of glycosylation does not lead to loss of antigen binding activity (DONADEL et al. 1994). Human SC purified from milk is heavily glycosylated with 5–7 N-linked sugar side chains, accounting for over 20% of its molecular mass (MIZOGUCHI et al. 1982). IL-4 and IL-5 have recently been shown to alter the terminal glycosylation pattern of IgA (CHINTALACHARUVU and EMANCIPATOR 1997). These authors suggest that increased production of IL-4 and IL-5 by peripheral blood lymphocytes from IgA nephropathy patients might result in the production of abnormally

glycosylated IgA which, in turn, may promote the disease by deposition of IgA in glomeruli.

### 5.1.3 Regulation of SC and $\alpha$ Chain Gene Expression

Poly-Ig receptor expression in mucosal epithelia is regulated by a complex interplay among lymphocytes, macrophages and epithelial cells. The presence of cytokine regulatory elements in the promoter of the poly-Ig receptor genes (PISKURICH et al. 1997) allows expression of the receptor to be upregulated in response to both Th1 and Th2 cytokines. Upregulation of poly-Ig receptor expression stimulates transcytosis of IgA antibodies. Several interferon-stimulated response elements (ISRE) have been identified in the human poly-Ig receptor promoter and first exon (PISKURICH et al. 1997; VERRIJDT et al. 1997). In HT-29 human colon carcinoma cells, poly-Ig receptor is upregulated by interferon gamma (IFN- $\gamma$ ) (PISKURICH et al. 1993), TNF- $\alpha$  (KVALE et al. 1988), and IL-4 (PHILLIPS et al. 1990). In contrast to the promoter ISREs, the ISRE in the first exon binds interferon responsive factor (IRF)-1, a member of the interferon regulatory-factor family, only after stimulation of HT-29 cells with IFN- $\gamma$ . Many microorganisms are able to stimulate cytokine or chemokine secretion directly by epithelial cells or indirectly via intraepithelial lymphocytes or inflammatory cells recruited into mucosal tissues (ISHIKAWA et al. 1993; QUIDING et al. 1991). Upregulation of poly-Ig receptor by cytokines involves a protein tyrosine kinase-dependent signaling pathway (DENNING 1996), in which the cytokine receptors recruit cytosolic tyrosine kinases (Janus kinases or JAKs) which, in turn, activate transcription factors of the STAT (signal transduction and activation of transcription) family (IHLE et al. 1994).

Several enhancer regions (DARIAVACH et al. 1991; MATTHIAS and BALTIMORE 1993) and hypersensitive sites (MADISEN and GROUDINE 1994) identified downstream of the murine C $\alpha$  gene are involved in its developmental and transcriptional regulation. The best characterized enhancer element, 3' $\alpha$ E, contains motifs (octamer,  $\kappa$ B, G-rich sequence) that mediate upregulation of transcription during maturation to plasma cells (SINGH and BIRSHTEN 1993; MEYER et al. 1995) as well as motifs (Elf-1, AP-1) for activation induced by B cell receptor cross-linking (GRANT et al. 1995). 3' $\alpha$ E also comprises the hypersensitive sites HS1 and HS2, which share 90% identity with their human  $\alpha$ 1 and  $\alpha$ 2 gene counterparts (MILLS et al. 1997). The HS1,2 core homology is likely to contain essential motifs important for the strong late B cell-specific enhancer characteristics of HS1,2 in mice and humans. Although the function of transcription binding sites in the human HS1,2 has not yet been demonstrated experimentally, this sequence element comprises Oct, AP-1, Ets,  $\mu$ E5 motifs (MILLS et al. 1997), all of which are nearly identical to, and functional in, the murine HS1,2 enhancer.

## 5.2 Functions of sIgA Antibodies

It is usually accepted that sIgA in secretions bathing the mucosal surfaces protect these surfaces against environmental pathogens by cross-linking microorganisms or macromolecules, thus facilitating their elimination by peristalsis or mucociliary movement and preventing their contact with the surface of epithelial cells, a phenomenon called immune exclusion. The molecular mechanisms underlying sIgA antibody-mediated protection still remain poorly understood, and it is likely that this unique effector molecule plays multiple roles in addition to immune exclusion (Fig. 2).

The ability of secretory antibodies to recognize intact bacteria, viruses or parasites at mucosal surfaces is a prerequisite for protection. This is reflected by the observation that polyclonal sIgA responses to mucosal pathogens is dominated by antibodies recognizing microbial surface antigens. An antibody response is not restricted to pathogens, but commensals such as gram-negative microorganisms introduced into the gut of germ-free mice also produce an immune response (SHROFF et al. 1995). This suggests that a successful sIgA response can attenuate chronic stimulation of germinal-center reactions in gut-associated lymphoid tissue in response to bacteria persisting in the gut. The role of sIgA in the establishment and maintenance of the gut flora will require further investigation.

It has been difficult to evaluate the relative importance of sIgA in protection or to assess whether "immune exclusion" by sIgA alone is able to prevent mucosal infection. Techniques for the production of monomeric IgA antibodies have been developed by several groups during the last few decades (STYLES et al. 1984; COLWELL et al. 1986; RITS et al. 1986; MAZANEC et al. 1987; WELTZIN et al. 1989), and production of recombinant sIgA has been achieved in mammalian cells (CHINTALACHARUVU and MORRISON 1997; BERDOZ et al. 1998) and in plants (MA et al. 1995). This has allowed assessment of protection in vivo either by passive transfer of monoclonal antibodies specific for viruses (MAZANEC et al. 1987; RENEGAR and SMALL 1991), bacteria (MICHETTI et al. 1992) or toxins (APTER et al. 1991; APTER et al. 1993b). The implantation of IgA-producing hybridoma cells into the backs of mice has also allowed efficient delivery of antibodies into mucosal sites (WINNER et al. 1991). The development of methods (CZERKINSKY et al. 1983; HANEBERG et al. 1994) to measure antibodies in mucosal tissues has also contributed to a better understanding of their role in protection.

### 5.2.1 Immune Exclusion by Antigen Cross-Linking

Resistance to mucosal infection has been correlated with specific sIgA. The secretory antibodies provide an immunological barrier that prevents foreign antigens, including bacteria, viruses, parasites and toxins, from attaching to mucosal surfaces (for review see NEUTRA et al. 1991). Due to their multivalency, sIgA antibodies are ideally designed to cross-link target macromolecules or microorganisms in the mucosal environment. They do so by preventing their diffusion through the glycocalyx at the surface of the epithelial cells, by blocking their binding to epithelial

surface receptors, by inhibiting their motility or by facilitating their entrapment in mucus. Prevention of viral attachment and subsequent internalization represents a major mechanism by which IgA-coated viruses are neutralized (TAYLOR and DIMMOCK 1985). In the absence of mucus and clearance mechanisms, sIgA antibodies are able to protect epithelial cell monolayers (MICHETTI et al. 1994). Aggregation of the pathogens, however, requires antibody concentrations which probably cannot be reached in the mucosal environment. This suggests that protection mediated by sIgA must involve other mechanisms.

### 5.2.2 Interaction with Mucus

Mucus is generally thought to protect epithelial cells by forming a diffusional barrier with pore size of about 100 nm, through which only small molecules including antibodies can pass. IgG, IgG fragments, IgA and IgM diffuse as rapidly in cervical mucus as in water, and it has been proposed that particles as large as viruses can similarly diffuse through human cervical mucus, provided that the particle forms no adhesive interactions with mucus glycoproteins (SALTZMAN et al. 1994). Cross-linking of viral particles by sIgA antibodies, however, significantly impedes their diffusion in cervical mucus. Whether mucus from other mucosal surfaces behave similarly remains to be established. Coating hydrophobic bacteria such as *Salmonella typhimurium* or *Escherichia coli* with sIgA renders their surface more hydrophilic, thus facilitating their entry and movement into mucus gels (MAGNUSSON and STJERNSTROEM 1982; MCCORMICK et al. 1988). Thus, microorganisms coated with sIgA antibodies would be more readily retained in the moving stream of mucus. Secretory antibodies would bind to mucus only when tightly packed on the surface of the microorganisms which increases the avidity of the low affinity IgA-mucin interaction.

### 5.2.3 Neutralization and Opsonization

Most IgA antibodies do not neutralize microorganisms by standard mechanisms used by IgG. Generally, they do not opsonize via the macrophage Fc $\gamma$  receptor in vitro, they do not activate the classical complement pathway and, thus, they do not lyse bacteria (KILIAN et al. 1988). There is evidence that sIgA can directly block the microbial sites that mediate epithelial attachment, either by binding to their adhesins or by sterically hindering their interaction with the epithelial cell surface (WILLIAMS and GIBBONS 1972; SVANBORG-EDEN and SVENNERHOLM 1978; WELTZIN et al. 1996). IgA antibodies which protect mucosal surfaces were shown to be inefficient when injected systemically (MICHETTI et al. 1992), suggesting that neutralization in a mucosal environment operates through different mechanisms than systemically (NEUTRA et al. 1994).

Several studies have demonstrated that protection of epithelial surfaces against viruses, bacteria or toxins can be accomplished with monoclonal IgA antibodies directed against microbial surface epitopes not necessarily involved in pathogenesis or adhesion (APTER et al. 1993a; PHALIPON et al. 1995).

Secretory IgA antibodies are able to cooperate with the innate defense system and enhance, for instance, the bacteriostatic activity of lactoperoxidase or lactoferrin present in mucosal fluids (TENOVUO et al. 1982). Inflammatory cells (neutrophils, eosinophils and macrophages) and immune cells (lymphocytes and monocytes) express Fc $\alpha$  receptors in mucosal tissues (FANGER et al. 1983; MONTEIRO et al. 1990). Occupancy and aggregation of the Fc $\alpha$  receptor on monocytes release pro-inflammatory cytokines (PATRY et al. 1995). The role of IgA-dependent cellular cytotoxicity which has been reported for the clearance of *Salmonella typhi* in mice (TAGLIABUE et al. 1985) has to be further investigated. It is likely that cell-mediated phenomena play an important role in protection of the host against mucosal and systemic infection once the pathogen has crossed the epithelial barrier and entered mucosal tissues.

#### 5.2.4 Intracellular Neutralization and Antigen Clearance

Microorganisms that are internalized by epithelial cells in mucosal tissues and accumulate in endosomes could be neutralized intracellularly in endosomal compartments provided that there is a meeting point between virus-containing endosomes and IgA-containing endosomal vesicles. This has been tested experimentally: IgA monoclonal antibodies against Sendai virus, a parainfluenza virus, co-localized with the viral hemagglutinin-neuraminidase protein within infected epithelial cells and reduced intracellular viral titers (MAZANEC et al. 1995).

A mouse model and "backpack tumor" transplantation (WINNER et al. 1991) were recently used to determine the protective effect of antibodies against rotavirus capsid proteins. Two non-neutralizing IgA antibodies to VP6 were capable of preventing virus shedding in infected mice (BURNS et al. 1996). These antibodies were not active when administered to the luminal side of the intestinal tract. The lack of virus shedding in the stools was taken as an indication that virus infection could be prevented inside the infected cells. However, the decrease of virus shedding could result from inhibition of virus assembly. It is difficult to imagine how VP6 specific antibodies that do not have access to surface-exposed epitopes on the virus could prevent infection. In addition, rotavirus infects villus epithelial cells, particularly those on the upper villus and IgA antibodies are transported mainly in crypt cells. Thus, it is not clear how IgA antibodies can neutralize the virus intracellularly. When prevention of diarrhea was assessed, rather than virus shedding, anti-VP6 antibodies were unable to prevent diarrhea in newborn mice (RUGGERI et al. 1998).

Soluble dimeric IgA-antigen immune complexes are transported from the basolateral to the apical surface of poly-Ig receptor-expressing epithelial monolayers and released into the apical compartment (KAETZEL et al. 1991; KAETZEL et al. 1994). This indicates that the pathways of epithelial transcytosis of free and complexed dimeric IgA are the same. Given the high population density of mucosal IgA plasma cells and the enormous surface area of poly-Ig receptor-expressing mucosal epithelium, it is likely that significant local transcytosis of IgA immune complexes occurs in vivo, providing a means to eliminate microorganisms that have

gained access to mucosal tissues. Such a mechanism may thus provide an important defense function for IgA.

*Acknowledgements.* We are grateful to the current and former members of our laboratories who have contributed to the work summarized in this review. We would like to thank Marian R. Neutra for stimulating discussion, and for critically reading the manuscript. The authors are supported by the Swiss National Science Foundation (Grants 31-47110-96 and 31-47296.96) and the Swiss Research Against Cancer (AKT 622).

## References

- Apodaca G, Katz LA, Mostov KE (1994) Receptor-mediated transcytosis of IgA in MDCK cells is via apical recycling endosomes. *J Cell Biol* 125:67-86
- Apter FM, Lencer WJ, Mekalanos JJ, Neutra MR (1991) Analysis of epithelial protection by monoclonal IgA antibodies directed against cholera toxin B subunits in vivo and in vitro. *J Cell Biol* 115:399a
- Apter FM, Lencer WJ, Finkelstein RA, Mekalanos JJ, Neutra MR (1993a) Monoclonal immunoglobulin A antibodies directed against cholera toxin prevent the toxin-induced chloride secretory response and block toxin binding to intestinal epithelial cells in vitro. *Infect Immun* 61:5271-5278
- Apter FM, Michetti P, Winner LSI, Mack JA, Mekalanos JJ, Neutra MR (1993b) Analysis of the roles of antilipopolysaccharide and anti-cholera toxin immunoglobulin A (IgA) antibodies in protection against *Vibrio cholerae* and cholera toxin by use of monoclonal IgA antibodies in vivo. *Infect Immun* 61:5279-5285
- Bakos MA, Kurosky A, and Goldblum RM (1991) Characterization of a critical binding site for human polymeric Ig on secretory component. *J Immunol* 147:3419-3426
- Banting G, Brake B, Braghetta P, Luzio JP, Stanley KK (1989) Intracellular targeting signals of polymeric immunoglobulin receptors are highly conserved between species. *FEBS Lett* 254:177-183
- Barroso M, Sztul ES (1994) Basolateral to apical transcytosis in polarized cells is indirect and involved BFA and trimeric G protein sensitive passage through the apical endosome. *J Cell Biol* 124:83-100
- Bendelac A, Fearon DT (1997) Innate immunity - innate pathways that control acquired immunity. *Curr Opin Immunol* 9:1-3
- Bjerke K, Brandtzaeg P (1990) Terminally differentiated human intestinal B cells. J chain expression of IgA and IgG subclass-producing immunocytes in the distal ileum compared with mesenteric and peripheral lymph nodes. *Clin Exp Immunol* 82:411-415
- Brandtzaeg P (1994) Distribution and characterization of mucosal immunoglobulin-producing cells. In: PL Ogra, J Mestecky, ME Lamm, W Strober, JR McGhee, J Bienenstock (eds) *Handbook of mucosal immunity*, Academic Press, New York, pp 251-259
- Brandtzaeg P, Prydz H (1984) Direct evidence for an integrated function of J chain and secretory component in epithelial transport of immunoglobulins. *Nature* 311:71-73
- Briskin M, Winsorhines D, Shyjan A, Cochran N, Bloom S, Wilson J, McEvoy LM, Butcher EC, Kassam N, Mackay CR, Newman W, Ringler DJ (1997) Human mucosal addressin cell adhesion molecule-1 is preferentially expressed in intestinal tract and associated lymphoid tissue. *Am J Pathol* 151:97-110
- Bromander AK, Ekman L, Kopf M, Nedrud JG, Lycke NY (1996) IL-6-deficient mice exhibit normal mucosal IgA responses to local immunizations and *Helicobacter felis* infection. *J Immunol* 156:429-497
- Burns JW, Siadatpajouh M, Krishnaney AA, Greenberg HB (1996) Protective effect of rotavirus VP6-specific IgA monoclonal antibodies that lack neutralizing activity. *Science* 272:104-107
- Butcher EC, Picker LJ (1996) Lymphocyte homing and homeostasis. *Science* 272:60-66
- Cattaneo A, Neuberger MS (1987) Polymeric immunoglobulin M is secreted by transfectants of non-lymphoid cells in the absence of immunoglobulin J chain. *EMBO J* 6:27532-758
- Cebra JJ, Logan AC, Weinstein PD (1991) The preference for switching to expression of the IgA isotype of antibody exhibited by B lymphocytes in Peyer's patches is likely due to intrinsic properties of their microenvironment. *Immunol Res* 10:393-395
- Chapin SJ, Enrich C, Aroeti B, Havel RJ, Mostov KE (1996) Calmodulin binds to the basolateral targeting signal of the polymeric immunoglobulin receptor. *J Biol Chem* 271:333-342

- important
- who have  
leutra for  
the Swiss  
h Against
- cells is via
- onoclonal  
15:399a  
oglobulin  
onse and
- le roles of  
rotection  
ct Immun
- or human
- s of poly-  
-183  
involved  
4:83-100  
ity. Curr
- ression of  
teric and
- cells. In:  
f mucosal
- secretory
- , Kassam  
ecule-I is  
:97-110  
it normal  
156:429
- rus VP6-
- s of non-
- A isotype  
s of their
- isolateral
- Chintalacharuvu KR, Morrison SL (1997) Production of secretory immunoglobulin A by a single mammalian cell. *Proc Natl Acad Sci U S A* 94:366-368
- Chintalacharuvu SR, Emancipator SN (1997) The glycosylation of IgA produced by murine B cells is altered by Th2 cytokines. *J Immunol* 159:232-233
- Colwell DE, Michalek SM, McGhee JR (1986) Method for generating a high frequency of hybridomas producing monoclonal IgA antibodies. *Methods Enzymol* 121:42-51
- Corthésy B, Kaufmann M, Phalipon A, Peitsch MC, Neutra MR, Kraehenbuhl J-P (1996) A pathogen-specific epitope inserted into recombinant secretory immunoglobulin A is immunogenic by the oral route. *J Biol Chem* 271:33670-33677
- Coyne RS, Siebrecht M, Peitsch MC, Casanova JE (1994) Mutational analysis of polymeric immunoglobulin receptor/ligand interactions. *J Biol Chem* 269:31620-31625
- Czerkinsky C, Nilsson LA, Nygren H, Ouchterlony O, Tarkowski A (1983) A solid-phase enzyme-linked immunospot (ELISPOT) assay for enumeration of specific antibody-secreting cells. *J Immunol Methods* 65:109-121
- Dariavach P, Williams GT, Campbell K, Pettersson S, Neuberger MS (1991) The mouse IgH 3'-enhancer. *Eur J Immunol* 21:149-1504
- Davis AC, Roux KH, Pursey J, Shulman MJ (1989) Intermolecular disulfide bonding in IgM: effects of replacing cysteine residues in the mu heavy chain. *EMBO J* 8:521-526
- Defrance T, Vanbervliet B, Briere F, Durand I, Rousset F, Banchereau J (1992) Interleukin 10 and transforming growth factor beta cooperate to induce anti-CD40-activated naive human B cells to secrete immunoglobulin A. *J Exp Med* 175:67-82
- Deitcher DL, Mostov KE (1986) Alternate splicing of rabbit polymeric immunoglobulin receptor. *Mol Cell Biol* 6:2712-2715
- Denning GM (1996) IL-4 and IFN-gamma synergistically increase total polymeric IgA receptor levels in human intestinal epithelial cells. Role of protein tyrosine kinases. *J Immunol* 156:480-484
- Donadel G, Calabro A, Sigounas G, Hascall VC, Notkins AL, Harindranath N (1994) Human polyreactive and monoreactive antibodies: effect of glycosylation on antigen binding. *Glycobiology* 4:49-96
- Eilfert H, Quentin E, Decker J, Hillemeir S, Hufschmidt M, Klingmüller D, Weber MH, Hilschmann N (1984) Die Primärstruktur der menschlichen Sekretkomponente und die Anordnung der Disulfidbrücken. *Hoppe-Seyler's Z Physiol Chem* 365:1489-1495
- Endo T, Mestecky J, Kulhavy R, Kobata A (1994) Carbohydrate heterogeneity of human myeloma proteins of the IgA1 and IgA2 subclasses. *Mol Immunol* 31:141-142
- Fanger MW, Goldstine SN, Shen L (1983) The properties and role of receptors for IgA on human leukocytes. *Ann N Y Acad Sci* 409:552-563
- Fearon DT, Locksley RM (1996) Elements of immunity - the instructive role of innate immunity in the acquired immune response. *Science* 272:50-54
- Förster R, Mattis AE, Kremmer E, Wolf E, Brem G, Lipp M (1996) A putative chemokine receptor, BLR1, directs B cell migration to defined lymphoid organs and specific anatomic compartments of the spleen. *Cell* 87:1037-1047
- Frutiger S, Hughes GJ, Fonck C, Jaton JC (1987) High and low molecular weight rabbit secretory components. Evidence for the deletion of the second and third domains in the smaller polypeptide. *J Biol Chem* 262:1712-1715
- Grant PA, Thompson CB, Pettersson S (1995) IgM receptor-mediated transactivation of the IgH 3' enhancer couples a novel Elf-1-AP-1 protein complex to the developmental control of enhancer function. *EMBO J* 14:4501-4513
- Gunn MD, Ngo VN, Ansel KM, Ekland EH, Cyster JG, Williams LT (1998) A B-cell-homing chemokine made in lymphoid follicles activates Burkitts-lymphoma receptor-1. *Nature* 391:799-803
- Haneberg B, Kendall D, Amerongen HM, Apter FM, Kraehenbuhl J-P, Neutra MR (1994) Induction of specific immunoglobulin A in the small intestine, colon-rectum, and vagina measured by a new method for collection of secretions from local mucosal surfaces. *Infect Immun* 62:15-23
- Hayami K, Fukuta D, Nishikawa Y, Yamashita Y, Inui M, Ohyama Y, Hikida M, Ohmori H, Takai T (1997) Molecular cloning of a novel murine cell-surface glycoprotein homologous to killer cell inhibitory receptors. *J Biol Chem* 272:7320-7327
- Hendrickson BA, Conner DA, Ladd DJ, Kendall D, Casanova JE, Corthésy B, Max EE, Neutra MR, Seidman CE, Seidman JG (1995) Altered hepatic transport of immunoglobulin A in mice lacking the J chain. *J Exp Med* 182:1905-1911
- Hendrickson BA, Rindisbacher L, Corthésy B, Kendall D, Waltz DA, Neutra MR, Seidman JG (1996) Lack of association of secretory component with IgA in J chain-deficient mice. *J Immunol* 157:750-754



- Hirt RP, Hughes GJ, Frutiger S, Michetti P, Perregaux C, Jeanguenat N, Neutra MR, Kraehenbuhl J-P (1993) Transcytosis of the polymeric Ig receptor requires phosphorylation of serine 664 in the absence but not the presence of dimeric IgA. *Cell* 74:245-255
- Hunkapiller T, Hood L (1989) Diversity of the immunoglobulin gene superfamily. *Adv Immunol* 44:1-63
- Ihle JN, Witthuhn BA, Quelle FW, Yamamoto K, Thierfelder WE, Kreider B, Silvennoinen O (1994) Signaling by the cytokine receptor superfamily: JAKs and STATs. *Trends Biochem Sci* 19:222-227
- Ishikawa H, Li Y, Abeliovich A, Yamamoto S, Kaufmann SH, Tonegawa S (1993) Cytotoxic and interferon gamma-producing activities of gamma delta T cells in the mouse intestinal epithelium are strain dependent. *Proc Natl Acad Sci U S A* 90:820-208
- Israel EJ, Taylor S, Wu Z, Mizoguchi E, Blumberg RS, Bhan A, Simister NE (1997) Expression of the neonatal Fc receptor, FcRn, on human intestinal epithelial cells. *Immunology* 92:69-74
- Kaetzel CS, Robinson JK, Chintalacharuvu KR, Vaerman J-P, Lamm M (1991) The polymeric immunoglobulin receptor (secretory component) mediates transport of immune complexes across epithelial cells: a local defense function for IgA. *Proc Natl Acad Sci U S A* 88:8796-8800
- Kaetzel CS, Robinson JK, Lamm ME (1994) Epithelial transcytosis of monomeric IgA and IgG cross-linked through antigen to polymeric IgA. A role for monomeric antibodies in the mucosal immune system. *J Immunol* 152:72-76
- Kawamura S, Saitou N, Ueda S (1992) Concerted evolution of the primate immunoglobulin alpha-gene through gene conversion. *J Biol Chem* 267:353-367
- Kawanishi H, Saltzman LE, Strober W (1983) Mechanisms regulating IgA class-specific immunoglobulin production in murine gut-associated lymphoid tissues. I. T cells derived from Peyer's patches that switch. *J Exp Med* 157:433-450
- Kerr MA (1990) The structure and function of human IgA. *Biochem J* 271:285-296
- Kilian M, Mestecky J, Russell MW (1988) Defense mechanisms involving Fc-dependent functions of immunoglobulin A and their subversion by bacterial immunoglobulin A proteases. *Microbiol Rev* 52:296-303
- Krajci P, Solberg R, Sandberg M, Oyen O, Jahnsen T, Brandtzaeg P (1989) Molecular cloning of the human transmembrane secretory component (poly-Ig receptor) and its mRNA expression in human tissues. *Biochem Biophys Res Com* 158:783-789
- Kvale D, Lovhaug D, Sollid LM, Brandtzaeg P (1988) Tumor necrosis factor-alpha upregulates expression of secretory component, the epithelial receptor for polymeric Ig. *J Immunol* 140:3086-3089
- Legler DF, Loetscher M, Stuber Roos R, Clark-Lewis I, Baggiolini M, Moser B (1998) B cell-attracting chemokine 1, a human CXC chemokine expressed in lymphoid tissues, selectively attracts B lymphocytes via BRL1/CXCR5. *J Exp Med* 187:665-660
- Li MJ, Peakman MC, Golub EI, Reddy G, Ward DC, Radding CM, Maizels N (1996) Rad51 expression and localization in B cells carrying out class switch recombination. *Proc Natl Acad Sci USA* 93:1022-1027
- Liu YJ, Mason DY, Johnson GD, Abbot SD, Gregory CD, Hardie DL, Gordon J, MacLennan IC (1991) Germinal center cells express bcl-2 protein after activation by signals which prevent their entry into apoptosis. *Eur J Immunol* 21:1905-1910
- Liu YJ, Johnson GD, Gordon J, MacLennan ICM (1992) Germinal centres in T-cell-dependent antibody responses. *Immunol Today* 13:17-21
- Lüllau E, Heyse S, Vogel H, Marison I, von Stockar U, Kraehenbuhl J-P, Corthesy B (1996) Antigen binding properties of purified IgA and reconstituted secretory IgA antibodies. *J Biol Chem* 271:16300-16309
- Lycke N, Strober W (1989) Cholera toxin promotes B cell isotype differentiation. *J Immunol* 142:3781-3787
- Ma JKC, Hiatt A, Hein M, Vine ND, Wang F, Stabila P, Vandolleweerd C, Mostov K, Lehner T (1995) Generation and assembly of secretory antibodies in plants. *Science* 268:716-719
- Madisen L, Groudine M (1994) Identification of a locus control region in the immunoglobulin heavy-chain locus that deregulates c-myc expression in plasmacytoma and Burkitt's lymphoma cells. *Genes Dev* 8:2212-2226
- Magnusson KE, Stjernstroem I (1982) Mucosal barrier mechanisms. Interplay between secretory IgA (sIgA), IgG and mucins on the surface properties and association of salmonellae with intestine and granulocytes. *Immunology* 45:239-248
- Matsuda H, Watanabe N, Kiso Y, Hirota S, Ushio H, Kannan Y, Azuma M, Koyama H, Kitamura Y (1990) Necessity of IgE antibodies and mast cells for manifestation of resistance against larval *Haemaphysalis longicornis* ticks in mice. *J Immunol* 144:25-62

- nbuhl J-P  
e absence
- l 44:1-63  
O (1994)  
:222-227  
toxic and  
elium are
- on of the
- ic immu-  
epithelial
- gG cross-  
l immune
- pha-gene
- oglobulin  
ches that
- ctions of  
biol Rev
- ng of the  
n human
- regulates  
40:3086-
- attracting  
s B lym-
- xpression  
93:1022-
- IC (1991)  
ntry into
- antibody
- Antigen  
l Chem
- 42:3781-
- T (1995)
- n heavy-  
ls. Genes
- tory IgA  
stine and
- amura Y  
rval Ha-
- Matsumoto R, Matsumoto M, Mita S, Hitoshi Y, Adno M, Araki S, Yamaguchi N, Tominaga A, Takatsu K (1989) Interleukin-5 induces maturation but not class switching of surface IgA-positive B cells into IgA-secreting cells. *Immunology* 66:32-41
- Matthias P, Baltimore D (1993) The immunoglobulin heavy chain locus contains another B-cell-specific 3' enhancer close to the alpha constant region. *Mol Cell Biol* 13:154-155
- Mazanec MB, Nedrud JG, Lamm ME (1987) Immunoglobulin A monoclonal antibodies protect against Sendai virus IgA nephropathy: pathogenesis of the most common form of glomerulonephritis. *J Virol* 61:2624-2626
- Mazanec MB, Kaetzel CS, Lamm ME, Fletcher D, Peterra J, Nedrud JG (1995) Intracellular neutralization of Sendai and influenza viruses by IgA monoclonal antibodies. *Adv Exp Med Biol* 371A:651-654
- McCormick BA, Stocker BA, Laux DC, Cohen PS (1988) Roles of motility, chemotaxis, and penetration through and growth in intestinal mucus in the ability of an avirulent strain of *Salmonella typhimurium* to colonize the large intestine of streptomycin-treated mice. *Infect Immun* 56:220-217
- McGhee JR, Fujihashi K, Beagley KW, Kiyono H (1991) Role of interleukin-6 in human and mouse mucosal IgA plasma cell responses. *Immunol Res* 10:418-422
- Mestecky J, McGhee JR (1987) Immunoglobulin A (IgA): molecular and cellular interactions involved in IgA biosynthesis and immune response. *Adv Immunol* 40:153-245
- Mestecky J, Lue C, Russell MW (1991) Selective transport of IgA: cellular and molecular aspects. *Gastroenterol Clin North Am* 20:441-471
- Meyer KB, Skogberg M, Margenfeld C, Ireland J, Pettersson S (1995) Repression of the immunoglobulin heavy chain 3' enhancer by helix-loop-helix protein Id3 via a functionally important E47/E12 binding site: implications for developmental control of enhancer function. *Eur J Immunol* 25:1770-1777
- Michetti P, Mahan MJ, Slauch JM, Mekalanos JJ, Neutra MR (1992) Monoclonal secretory immunoglobulin A protects mice against oral challenge with the invasive pathogen *Salmonella typhimurium*. *Infect Immun* 60:1786-1792
- Michetti P, Porta N, Mahan MJ, Slauch JM, Mekalanos JJ, Blum AL, Kraehenbuhl J-P, Neutra MR (1994) Monoclonal immunoglobulin A prevents invasion of polarized epithelial cell monolayers by *Salmonella typhimurium*. *Gastroenterology* 107:915-923
- Mills FC, Harindranath N, Mitchell M, Max EE (1997) Enhancer complexes located downstream of both human immunoglobulin C alpha genes. *J Exp Med* 186:84-88
- Mizoguchi A, Mizuochi T, Kobata A (1982) Structures of the carbohydrate moieties of secretory component purified from human milk. *J Biol Chem* 257:961-961
- Monteiro RC, Kubagawa H, Cooper MD (1990) Cellular distribution, regulation, and biochemical nature of an Fc alpha receptor in humans. *J Exp Med* 171:597-613
- Mostov KE, Deitcher DL (1986) Polymeric immunoglobulin receptor expressed in MDCK cells transcytoses IgA. *Cell* 46:613-621
- Mostov KE, Friedlander M, Blobel G (1984) The receptor for transepithelial transport of IgA and IgM contains multiple immunoglobulin-like domains. *Nature* 308:37-43
- Mostov KE, Apodaca G, Aroeti B, Okamoto C (1992) Plasma membrane protein sorting in polarized epithelial cells. *J Cell Biol* 116:577-583
- Natvig IB, Johansen FE, Nardeng TW, Haranldsen G, Brandtzag P (1997) Mechanism for enhanced external transfer of dimeric IgA over pentameric IgM. *J Immunol* 159:4330-4340
- Nedrud JG, Blanchard T, Czinn S, Harriman GR (1996) Orally immunized IgA deficient mice are protected against *H. felis* infection. *Gut* 39 [Suppl 2]:A-45
- Neutra MR, Weltzin R, Winner L, Mack J, Michetti P, Morrison L, Fields BN, Mekalanos JJ, Kraehenbuhl J-P (1991) Identification and use of protective monoclonal IgA antibodies against viral and bacterial pathogens. *Adv Exp Med Biol* 310:179-182
- Neutra MR, Michetti P, Kraehenbuhl J-P (1994) Secretory immunoglobulin A: induction, biogenesis and function. In: L R Johnson (ed) *Physiology of the gastrointestinal tract*. Raven, New York, pp 685-709
- Patry C, Herbelin A, Lehen A, Bach JF, Monteiro RC (1995) Fc alpha receptors mediate release of tumour necrosis factor-alpha and interleukin-6 by human monocytes following receptor aggregation. *Immunology* 86:1-5
- Phalipon A, Kaufmann M, Michetti P, Cavaillon JM, Huerre M, Sansonetti PJ, Kraehenbuhl J-P (1995) Monoclonal IgA antibody directed against serotype-specific epitope of *Shigella flexneri* liposaccharide protects against murine experimental shigellosis. *J Exp Med* 182:769-778

- Phillips JO, Everson MP, Moldoveanu Z, Lue C, Mestecky J (1990) Synergistic effect of IL-4 and IFN-gamma on the expression of polymeric Ig receptor (secretory component) and IgA binding by human epithelial cells. *J Immunol* 145:174-174
- Piskurich JF, France JA, Tamer CM, Willmer CA, Kaetzel CS, Kaetzel DM (1993) Interferon-gamma induces polymeric immunoglobulin receptor mRNA in human intestinal epithelial cells by a protein synthesis dependent mechanism. *Mol Immunol* 30:413-421
- Piskurich JF, Yougman KR, Phillips KM, Hempen PM, Blanchard MH, France JA, Kaetzel CS (1997) Transcriptional regulation of the human polymeric immunoglobulin receptor gene by interferon-gamma. *Mol Immunol* 34:75-91
- Plaut AG (1988) Production and isolation of neisserial IgA proteases. *Methods Enzymol* 165:11-20
- Plaut AG, Gilbert JV, Leger G, Blumenstein M (1985) IgA1 protease cleaves heavy chains independently in dimeric human IgA1. *Mol Immunol* 22:821-826
- Plebani A, Mira E, Mevio E, Monafò V, Notarangelo LD, Avanzini A, Ugazio AG (1983) IgM and IgD concentrations in the serum and secretions of children with selective IgA deficiency. *Clin Exp Immunol* 53:689-696
- Quiding M, Nordstrom I, Kilander A, Andersson G, Hanson LA, Holmgren J, Czerkinsky C (1991) Intestinal immune responses in humans. Oral cholera vaccination induces strong intestinal antibody responses and interferon-gamma production and evokes local immunological memory. *J Clin Invest* 88:143-148
- Quiding-Jarbrink M, Nordstrom I, Granstrom G, Kilander A, Jertborn M, Butcher EC, Lazarovits AI, Holmgren J, Czerkinsky C (1997) Differential expression of tissue-specific adhesion molecules on human circulating antibody-forming cells after systemic, enteric, and nasal immunizations. A molecular basis for the compartmentalization of effector B cell responses. *J Clin Invest* 99:128-286
- Ramsay AJ, Husband AJ, Ramshaw IA, Bao S, Matthaei KI, Koehler G, Kopf M (1994) The role of interleukin-6 in mucosal IgA antibody responses in vivo. *Science* 264:561-563
- Renegar KB, Small PA (1991) Passive transfer of local immunity to influenza virus infection by IgA antibody. *J Immunol* 146:1972-1978
- Rindischbacher L, Cotter S, Wittek R, Kraehenbuhl J-P, Corthesy B (1995) Production of human secretory component with dimeric IgA binding capacity using viral expression systems. *J Biol Chem* 270:14220-14228
- Rits M, Cormont F, Bazin H, Maykens R, Vaerman J-P (1986) Rat monoclonal antibodies. VI. Production of IgA secreting hybridomas with specificity for the 2,4-dinitrophenyl (DNP) hapten. *J Immunol Methods* 89:81-87
- Roberts DM, Guenther M, Rodewald R (1990) Isolation and characterization of the Fc receptor from the fetal yolk sac of the rat. *J Cell Biol* 111:186-187
- Rodewald RD, Kraehenbuhl J-P (1984) Receptor-mediated transport of IgG. *J Cell Biol* 99:159-164
- Roque-Barreira MC, Campos-Neto A (1985) Jacalin: an IgA-binding lectin. *J Immunol* 134:1740-1743
- Rott LS, Rose JR, Bass D, Williams MB, Greenberg HB, Butcher EC (1997) Expression of mucosal homing receptor alpha-4-beta-7 by circulating Cd4(+) cells with memory for intestinal rotavirus. *J Clin Invest* 100:1204-1208
- Ruggeri FM, Johansen K, Basile G, Kraehenbuhl J-P, Svensson L (1998) Rotavirus IgA neutralize virus in vitro after transcytosis through epithelial cells and protect infant mice from diarrhea. *J Virol* (in press)
- Sallusto F, Mackay CR, Lanzavecchia A (1997) Selective expression of the cotaxin receptor CCR3 by human T helper 2 cells. *Science* 277:2005-2007
- Saltzman WM, Radomsky ML, Whaley KJ, Conc RA (1994) Antibody diffusion in human cervical mucus. *Biophysical J* 66:508-515
- Schaerer E, Neutra MR, Kraehenbuhl J-P (1991) Molecular and cellular mechanisms involved in transepithelial transport. *J Membr Biol* 123:93-103
- Schiff JM, Fisher MM, Jones AL, Underdown BJ (1986) Human IgA as a heterovalent ligand: switching from the asialoglycoprotein receptor to secretory component during transport across the rat hepatocyte. *J Cell Biol* 102:920-931
- Shroff KE, Meslin K, Cebra JJ (1995) Commensal enteric bacteria engender a self-limiting humoral mucosal immune response while permanently colonizing the gut. *Infect Immun* 63:3904-3913
- Singh M, Birshtein BK (1993) NF-HB (BSAP) is a repressor of the murine immunoglobulin heavy-chain 3' alpha enhancer at early stages of B-cell differentiation. *Mol Cell Biol* 13:3611-3622
- Solari R, Kraehenbuhl J-P (1984) Biosynthesis of the IgA antibody receptor: a model for the transepithelial sorting of a membrane glycoprotein. *Cell* 36:61-71

- and IFN- $\gamma$  by human
- on-gamma a protein
- CS (1997) interferon-
- 11 20 pendently
- I and IgD Exp Im-
- C (1991) antibody lin Invest
- azarovits molecules ations. A t 99:128-
- ae role of
- n by IgA
- secretory 70:14220-
- VI. Pro- hapten.
- stor from
- 59-164 740-1743 mucosal rotavirus.
- ilize virus Virol (in
- CCR3 by
- 1 cervical
- olved in
- switching e rat he-
- humoral 13 ivy-chain
- transepi-
- Solari R, Kraehenbuhl J-P (1987) Receptor-mediated transepithelial transport of polymeric immunoglobulins. In: MC Neville, CW Daniel (eds) *The mammary gland*. Plenum, New York, pp 269-298
- Song W, Apodaca G, Mostov K (1994) Transcytosis of the polymeric immunoglobulin receptor is regulated in multiple intracellular compartments. *J Biol Chem* 269:29474-29480
- Spieker-Polet H, Yam PC, Knight KL (1993) Differential expression of 13 IgA-heavy chain genes in rabbit lymphoid tissues. *J Immunol* 150:5457-5465
- Strober W, Ehrhardt OR (1994) Regulation of IgA B cell development. In: PL Ogra, J Mestecky, ML Lamm, W Strober, JR McGhee, J Bienenstock (eds) *Handbook of mucosal immunology*. Academic Press, New York, pp 159-176
- Styles JM, Dean CJ, Gyure A, Hobbs SM, Halé JG (1984) The production of hybridomas from the gut-associated lymphoid tissue of tumor-bearing rats. II. Peripheral intestinal lymph as a source of IgA producing cells. *Clin Exp Immunol* 57:365-370
- Svanborg-Eden C, Svennerholm AM (1978) Secretory immunoglobulin A and G antibodies prevent adhesion of *Escherichia coli* to human urinary tract epithelial cells. *Infect Immun* 22:790-797
- Tagliabue A, Villa L, Boraschi D, Peri G, de Gori V, Nencioni L (1985) Natural anti-bacterial activity against *Salmonella typhi* by human T4+ lymphocytes armed with IgA antibodies. *J Immunol* 135:4178-4182
- Taylor AK, Wall R (1988) Selective removal of alpha heavy-chain glycosylation sites causes immunoglobulin A degradation and reduced secretion. *Mol Cell Biol* 8:419-203
- Taylor HP, Dimmock NJ (1985) Mechanism of neutralization of influenza virus by secretory IgA is different from that of monomeric IgA or IgG. *J Exp Med* 161:198-209
- Tenovuio J, Moldoveanu Z, Mestecky J, Pruitt KM, Rahemtulla BM (1982) Interaction of specific and innate factors of immunity: IgA enhances the antimicrobial effect of the lactoperoxidase system against *Streptococcus mutans*. *J Immunol* 128:726-731
- Verrijdt G, Swinnen J, Peeters B, Verhoeven G, Rombauts W, Claessens F (1997) Characterization of the human secretory component gene promoter. *Biochim Biophys Acta* 1350:147-154
- Weimbs T, Low SH, Chapin SJ, Mostov KE (1997) Apical targeting in polarized epithelial cells - there's more afloat than rafts. *Trends Cell Biol* 7:393-399
- Weltzin RA, Lucia Jandris P, Michetti P, Fields BN, Kraehenbuhl J-P, Neutra MR (1989) Binding and transepithelial transport of immunoglobulins by intestinal M cells: demonstration using monoclonal IgA antibodies against enteric viral proteins. *J Cell Biol* 108:1673-1685
- Weltzin R, Traina-Dorge V, Soike K, Zhang JY, Mack P, Soman G, Drabik G, Monath TP (1996) Intranasal monoclonal IgA antibody to respiratory syncytial virus protects rhesus monkeys against upper and lower respiratory tract infection. *J Infect Dis* 174:256-261
- Williams RC, Gibbons RJ (1972) Inhibition of bacterial adherence by secretory immunoglobulin A: a mechanism of antigen disposal. *Science* 177:697-699
- Winner LSI, Mack J, Weltzin RA, Mekalanos JJ, Kraehenbuhl J-P, Neutra MR (1991) New model for analysis of mucosal immunity: intestinal secretion of specific monoclonal immunoglobulin A from hybridoma tumors protects against *Vibrio Cholerae* infection. *Infect Immun* 59:977-982
- Wold AE, Motas C, Svanborg C, Mestecky J (1994) Lectin receptors on IgA isotypes. *Scand J Immunol* 39:195-201
- Wold AE, Motas C, Svanborg C, Hanson LA, Mestecky J (1995) Characterization of IgA1, IgA2 and secretory IgA carbohydrate chains using plant lectins *Adv Exp Med Biol* 371A:585-589